

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Antibiotic resistance profiles in relation to virulence factors and phylogenetic groups of uropathogenic Escherichia coli isolated from dogs and cats

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/148721> since

Published version:

DOI:10.1007/s11259-011-9500-z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Abstracts

LXIV Meeting of the Italian Society for Veterinary Science (SISVet), Asti 2010 – selected papers

Full papers available in Pugliese A, Gaiti A, Boiti C (2012) *Veterinary Science – Current Aspects in Biology, Animal Pathology, Clinic and Food Hygiene*. Springer-Verlag, Heidelberg

Alberto Gaiti ¹ 

(1) Faculty of Veterinary Medicine, University of Perugia, Perugia, Italy

 Alberto Gaiti
Email: gaiti-a@unipg.it

Published online: 4 November 2011

ANTIBIOTIC-RESISTANCE PROFILES IN RELATION TO VIRULENCE FACTORS, AND PHYLOGENETIC GROUP OF UROPATHOGENIC *ESCHERICHIA COLI* ISOLATED FROM DOGS AND CATS

Tramuta C¹, Nucera D², Robino P¹, Salvarani S¹, Nebbia P¹ ¹*Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia;* ²*Dipartimento di Patologia Animale - Università di Torino*

SUMMARY - In this study we assessed the association among antibiotic resistance profiles, virulence genotype and phylogenetic group within a collection of *Escherichia coli*. Forty uropathogenic *E. coli* strains isolated from dogs and cats with urinary tract infection were analysed by disk diffusion method and by PCRs. Results suggest that clinical isolates with a number ≥ 3 of virulence factors belong on the whole to phylogroup B2 (90.5%) and *E. coli* strains with a number < 3 of virulence genes were distributed in all phylogenetic groups. Considering virulence factors association with antimicrobial resistance, no statistically significant results were obtained ($P > 0.05$). Only gene *iutA* showed a trend of association with MDR ($P = 0.055$). Resistant strains were distributed in all phylogenetic groups (57%, B2; 43%, non-B2), in contrast *E. coli* isolates susceptible were associated with group B2 (90%) and with group D (10%).