

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Development of an ELISPOT assay for HSV-1 and clinical validation in lung transplant patients

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1680861> since 2018-11-07T13:38:13Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Development of an ELISPOT assay for HSV-1 and clinical validation in lung transplant**
2 **patients.**

3 ^{1,*}Cristina Costa, ¹Alessia Di Nauta, ¹Massimo Rittà, ¹Franca Sinesi, ¹Gabriele Bianco, ¹Francesca
4 Sidoti, ²Paolo Solidoro, ¹Rossana Cavallo.

5 ¹Microbiology and Virology Unit, Laboratory of Virology; ²Division of Pneumology, Azienda
6 Ospedaliero Universitaria Città della Salute e della Scienza di Torino, Corso Bramante 88 – 10126
7 Turin, Italy.

8

9 **Running title.** Elispot for HSV-1.

10

11 ***Corresponding author:**

12 Dr. Cristina Costa

13 Microbiology and Virology Unit

14 University Hospital Città della Salute e della Scienza di Torino

15 Corso Bramante 88 – 10126 Turin, Italy

16 Phone: +39(11)6335953 Fax: +39(11)6335194

17 E-mail: cristina.costa@unito.it

18

19

20 **SUMMARY**

21 Cellular immunity plays a relevant role in control of HSV-1 infection/reactivation with a potential
22 impact on clinical-therapeutic management of immunocompromised patients, such as transplant
23 recipients.

24 Herein, we quantitatively evaluated T-cell response directed at HSV-1 by a newly developed IFN- γ
25 EliSPOT assay in 53 patients (including 45 lung transplant recipients and eight subjects in waiting
26 list).

27 Overall, 62.2% of transplant patients and 62.5% of subjects in waiting list evidenced a response to
28 HSV-1 with no significant difference in the level of virus-specific cellular immunity. Response
29 tended to be lower in the first 3 months posttransplantation with progressive recovery of
30 pretransplantation status by the second year and in the presence of HSV-1 DNA positivity in
31 bronchoalveolar lavage. As expected, no response was found in seronegative patients. No
32 significant difference in the level of response according to IgM and IgG status was found.

33 Further studies are required to define the role of HSV-1 specific immune response for the clinical-
34 therapeutic management of lung transplant patients and in other clinical settings and to define cut-
35 off levels discriminating between absence/low and strong response to be related to the risk of viral
36 infection/reactivation.

37

38 **Key words:** herpes simplex virus type 1; cellular immune response; EliSPOT assay; lung
39 transplantation.

40 **Introduction**

41 Herpes simplex virus type-1 (HSV-1) is a highly seroprevalent and ubiquitously distributed dsDNA
42 virus belonging to the *Herpesviridae* family, α -herpesvirinae subfamily. Primary infection usually
43 occurs early in the childhood and is followed by a lifelong latent infection in neurons of central
44 ganglia, from which reactivation may occur. Whereas asymptomatic mucosal shedding is common
45 and HSV-1 has been isolated from the saliva of 1-5% of healthy subjects (Tsakris and Pitiriga
46 2011), reactivation has been reported particularly in immunosuppressed and critical patients in
47 which, beside classical presentation, visceral or disseminated disease can occur, including extensive
48 mucocutaneous involvement, hepatitis, meningoencephalitis, and pneumonitis (Tsakris and
49 Pitiriga 2011; Simmoons-Smit et al. 2006; Costa et al. 2012c; Wilk et al. 2013; Preiser et al. 2003;
50 Bonizzoli et al. 2016). As regards the lower respiratory tract, HSV-1 has been reported in 16 up to
51 32% of the cases (Bruynseels et al. 2003; van den Brink et al. 2004; Daubin et al. 2005; Luyt et al.
52 2007; Linssen et al. 2008) and has been increasingly associated to pulmonary diseases, with poor
53 outcome and high mortality rates (Costa et al. 2012c; Luyt et al. 2007; Linssen et al. 2008; Ong et
54 al. 2004; Engelmann et al. 2007; Gooskens et al. 2007; De Vos et al. 2009; Bouza et al. 2011;
55 Scheithauer et al. 2010). Adaptive immunity plays a pivotal role in uncomplicated recovery from
56 HSV infection, as evidenced by severe complications observed in immunocompromised
57 individuals, although the kinetics and specificity of HSV-specific T-cells during primary infection
58 are poorly unknown (Ouwendijk et al. 2013). After resolution of acute infection, memory T-cells
59 are detected at moderate levels in blood of immunocompetent subjects, with a poly-specific T-cell
60 response directed at distinct HSV-1 tegument and capsid proteins (Jing et al. 2012; Moss et al.
61 2012). Blood HSV-specific T-cells express high levels of cytolytic molecules and secrete IFN- γ
62 upon antigenic recall (Ouwendijk et al. 2013); higher levels of IFN- γ production are associated with
63 polyfunctionality of T-cells and better control of chronic viral infection (Merindol et al. 2012;
64 Harari et al. 2006). Moreover, HSV-1-specific T-cells localize to sites of primary and recurrent
65 infections, as well as latency sites, contributing to control viral latency and reactivation (Ouwendijk
66 et al. 2013; Khanna et al. 2003; Gebhardt et al. 2009; Ariotti et al. 2012). Quantitative evaluation of
67 HSV-1-specific T-cell response in blood compartment and the study of the relation between this and
68 ability of controlling local reactivation in the lung could be relevant for the clinical management of
69 immunocompromised patients at risk of severe pulmonary complications. At moment, no assay for
70 evaluation of cellular immune response to HSV-1 is available, as well as no data on its potential

71 impact on clinical/therapeutic management of infection/reactivation in different categories of
72 patients have been evaluated.

73 In the present study, we quantitatively evaluated T-cell responses directed at HSV-1 by an newly
74 developed IFN- γ EliSPOT assay in a susceptible population such as lung transplant recipients and
75 investigated the role of systemic virus-specific immunity in determining the risk of viral
76 reactivation in the lower respiratory tract.

77

78

79 **Materials and methods**

80 *Subjects and specimens*

81 Cellular immune response to HSV-1 was evaluated in an observational, longitudinal and
82 prospective study by IFN- γ EliSPOT assay on peripheral blood mononuclear cell (PBMC)
83 specimens from all lung transplant recipients admitted to the University Hospital “Città della Salute
84 e della Scienza di Torino”, Turin, Italy (Regione Piemonte Transplant Centre) over a two-year
85 period. The Lung Transplant Centre of the Piemonte region is the first in Italy for activity volume.
86 Overall, 53 patients (M/F, 33/12; mean age \pm standard deviation, 47.8 \pm 15.2 years; range, 16-69),
87 including 45 lung transplant recipients in the first two years posttransplantation and eight subjects
88 in waiting list were prospectively evaluated. In this study population, one (in subjects in waiting
89 list) or at least three (in transplant patients) PBMC specimens were collected, accounting for an
90 overall number of 168 samples (160 from transplant recipients, including 81 from 27 patients with
91 three evaluations, 64 from 16 patients with four evaluations, and 15 from three patients with five
92 evaluations; eight specimens from individuals in waiting list). A pre-transplant evaluation of HSV-1
93 cellular immune response was also obtained for all the patients, but three. Pretransplant serological
94 data for HSV-1/2 (IgG and IgM serostatus) were extrapolated from the local Transplant Registry
95 and were available for all patients, in particular five individuals IgG-negative and 48 IgG-positive,
96 with five subjects being IgM-positive. Baseline characteristics of the enrolled patients are reported
97 in Table 1. Moreover, 42 healthy seropositive individuals (IgG-positive, IgM-negative), including
98 38 without recurrent HSV-1 infection and four with at least one episode of HSV-1 infection (herpes
99 labialis) in the previous 12 months, were also studied by a single EliSPOT determination.

100 All subjects provided written informed consent and the study was approved by the institutional
101 review board. According to our lung transplant center’s practice, all patients received prophylaxis
102 for HSV consisting in administration of acyclovir (400 mg twice daily; to be reduced in case of
103 kidney failure or suspended in case of ganciclovir or valganciclovir treatment for CMV). In
104 addition, all patients received a universal, prolonged and combined anti-viral prophylaxis for CMV,
105 irrespective of serological matching donor/recipient, consisting in the administration of ganciclovir
106 or valganciclovir (450 mg twice daily) from day 21 posttransplantation for 3 weeks associated to
107 CMV-Ig (Cytotect Biotest) at days 1, 4, 8, 15, and 30 (1.5 ml/kg body weight) and monthly up to 2
108 years posttransplantation, according to local practice. Long-term immunosuppression was
109 maintained with tacrolimus or cyclosporine A (in patients with cystic fibrosis as underlying

110 disease), mycophenolate mofetil and prednisone (to be tapered or discontinued). Follow-up
111 surveillance bronchoscopies (with bronchoalveolar lavage [BAL] and transbronchial biopsy) were
112 scheduled at 1, 3, 6, 9, 12, 18, and 24 months posttransplantation, for the evaluation of rejection and
113 infections in the lower respiratory tract, as previously described (Costa et al. 2012a; Costa et al.
114 2008; Costa et al. 2011; Costa et al. 2012b). Therefore, virological data for HSV-1 were available
115 on BAL specimens concomitantly collected with samples for EliSPOT assay in all the cases. HSV-1
116 was evaluated on BAL specimens by real-time PCR using a commercially available kit (HSV-1
117 ELITe MGB® kit, ELITechGroup) following automated extraction with the Qiasymphony (Qiagen,
118 Hilden, Germany) instrument. Rapid shell vial isolation with indirect immunofluorescence for
119 HSV-1 was also performed, as previously described (Costa et al. 2007).

120

121 *IFN- γ EliSPOT assay*

122 HSV-1 antigenic stimulus consisted of a freeze-thaw/sonicated viral lysate prepared from expanded
123 long-term cultures of Vero cells (kidney epithelial cells from African green monkey, as previously
124 described (Terlizzi et al. 2009), infected with the Human herpesvirus 1 ATCC® VR-260™
125 [American Type Culture Collection, Manassas, VA, USA]). Aliquots of viral preparation were
126 stored at -80°C until use. For virus titration, 96-well plates at 60-80% confluence of Vero cells were
127 inoculated with 100 μ l of 10-fold diluted virus for TCID₅₀ assay, obtaining an end-point titer of
128 3.16×10^8 TCID₅₀/ml. Sonication included thawing of the virus in ice and 3 cycles at 20% intensity
129 for 30 seconds using the Sonopuls Ultraschall-Homogenisatoren instrument (Bandelin electronic
130 GmbH, Berlin, Germany). Subsequently, the virus underwent a through UV irradiation for
131 inactivation, with two cycles per transilluminator set at 1.2 J/cm². UV inactivation was carried out
132 also on the RPMI 1640-medium (Sigma-Aldrich, St. Louis, MO, USA), used for the EliSPOT assay
133 (see below). In order to ascertain the effective inactivation of the virus, a rapid shell vial culture
134 assay followed by indirect immunofluorescence was performed, as previously described (Costa et
135 al. 2007), and resulted negative (Figure 1). For antigenic stimulus, serial dilutions from 10⁶ up to
136 10³ of the inactivated virus, starting at 3.16×10^8 TCID₅₀/ml were used. Dose response curves were
137 performed with the lysate preparation to determine the amount of antigenic stimulus to use in the
138 IFN- γ EliSPOT assay: in particular, on PBMCs obtained from four healthy controls and two lung
139 transplant recipients.

140 Whole blood was collected directly into CPT Vacutainer tubes (BD, Franklin Lakes, NJ, USA) and
141 PBMCs were separated by density gradient sedimentation according to manufacturer instructions,
142 with minor modifications. Briefly, blood samples were centrifuged at 1800 g for 20 min at room
143 temperature. The resulting mononuclear cell fraction was washed twice with phosphate buffered
144 saline (PBS 1x, pH 7.4). Resulting PBMCs were cryopreserved in fetal calf serum (PAA
145 Laboratories GmbH, Pasching, Austria) with 10% dimethyl sulfoxide, placed into Nalgene
146 Cryovials (Nalge Nunc, Rochester, NY, USA) at -80°C for ≥ 24 h prior to transfer to liquid nitrogen
147 for long-term storage.

148 The IFN- γ EliSPOT assay was performed as described elsewhere (Costa et al. 2012b). Briefly,
149 PBMCs were thawed in RPMI-1640 (Sigma-Aldrich, St. Louis, MO, USA) supplemented with 10%
150 fetal calf serum and 1% l-glutamine, washed twice and rested for at least 4 h in complete RPMI-
151 1640 at 37°C, 5% CO₂, before assay. Subsequently, cell viability and count were assessed by trypan
152 blue staining in Burker's chamber to a final concentration of 2×10^6 cells/ml. Peripheral blood
153 mononuclear cells were plated at 2×10^5 cells/well onto a 96-well microplate pre-coated with anti-
154 human IFN- γ monoclonal antibody (EliSPOT Interferon- γ Basis Kit; AID, Strassberg, Germany)
155 and incubated with viral preparations, as described above. For negative and positive controls, cells
156 incubated with supplemented RPMI -1640 medium alone and with 1 μ g/ml phytohemagglutinin
157 mitogen (supplied by ELITechGroup, Milan, Italy) were used, respectively. Following a 18-20 h
158 incubation at 37°C, 5% CO₂, the microplates were washed 8-times with washing buffer and
159 incubated with biotinylated anti-human IFN- γ mAb at 1 μ g/ml in VP buffer at room temperature in
160 a wet chamber, in the dark, for 2 h. Subsequently, the microplates were washed 8-times with
161 washing buffer and incubated with streptavidin-horseradish peroxidase solution diluted 1:1000 in
162 blocking buffer. Following another washing step, as before, substrate solution
163 (tetramethylbenzidine) was added for colour development at room temperature in the dark for 12-15
164 min. The chromogenic reaction was stopped by extensive washing with tap water and microplates
165 were allowed to completely dry before analysis. Results were analyzed using a computer-assisted
166 system (AID EliSPOT Reader System, AID). Data were expressed as spot-forming units (SFU)/ $2 \times$
167 10^5 cells, with each spot representing a single cell that produces IFN- γ , calculated by subtracting the
168 mean of SFU obtained in unstimulated negative control from the mean SFU obtained in the antigen-
169 stimulated wells.

170 *Statistical analysis*

171 For descriptive statistics, data were expressed as raw number and percentage. For statistical
172 analysis, chi square, t test, and analysis of variance (ANOVA, followed by Bonferroni post-test)
173 were applied, as appropriate. A p value <0.05 was considered significant. Statistical analysis was
174 performed using GraphPad Prism version 5 (GraphPad Software, San Diego, USA).

175

176

177 **Results**

178 *Validation of the EliSPOT assay*

179 Based on dose-response curves on preliminary EliSPOT assays performed in triplicate on PBMCs
180 from four IgG positive healthy controls and two IgG-positive lung transplant patients, serial HSV-1
181 lysate dilutions at 3.16×10^4 and 3.16×10^3 TCID₅₀/mL were associated to more robust and
182 reproducible responses, even though not at statistical level ($p = n.s.$)(Figure 2). Therefore, these
183 dilutions were used as antigenic stimuli for subsequent HSV1-EliSPOT evaluations on specimens
184 from study population. The results evidenced that 3.16×10^3 TCID₅₀/mL HSV-1 stimulus was
185 associated to higher responses in comparison to 3.16×10^4 TCID₅₀/mL (mean SFU/2 x 10⁵ PBMCs
186 \pm SD; 3.765 ± 5.516 versus 2.662 ± 4.531 , $p=0.048$), when considering samples from
187 posttransplantation setting (peak value of response for each patient)(Figure 3).

188

189 *Clinical evaluation*

190 Overall, 28/45 (62.2%) transplant patients and 5/8 (62.5%) patients in waiting list evidenced a
191 positive response to HSV-1 lysate, with level of response ranging from 1 to 211 and from 8 to 53
192 SFU/2 x 10⁵ PBMCs, respectively. No significant difference of response was found between
193 samples from pre- and post-transplant patients, considering both all specimens (mean SFU/2 x 10⁵
194 PBMCs \pm SD, 8.9 ± 15.9 versus 7.2 ± 25.6 ; $p = 0.870$) and specimens with peak value from each
195 patient (10.4 ± 18.1 versus 5.15 ± 25.6 ; $p = 0.120$). Subsequently, as the main risk of HSV-1
196 infection is in the first period posttransplantation (up to 3 months, particularly the first 30 days), we
197 compared the degree of specific cellular response in specimens collected in the first 3 months
198 (overall, 10 specimens from as many patients) versus those collected at > 3 months and found no
199 significant differences, although mean values of SFU/2 x 10⁵ PBMCs tended to be lower in the early
200 period in comparison to >3 months (mean \pm SD, 2.7 ± 5.5 versus 7.4 ± 27.3 ; $p = 0.638$).

201 Among healthy seropositive individuals, 39/42 (83.3%) evidenced a positive response to HSV-1
202 lysate (37 with no episode of recent HSV-1 infection in the previous 12 months, two with two
203 episodes of herpes labialis), with level of response ranging from 8 to 36 SFU/2 x 10⁵ PBMCs; in
204 three subjects no response was found (one with no episode of HSV-1 infection and two with an
205 episode of herpes labialis in the previous 12 months). No significant difference of response was
206 found between individuals with and without HSV-1 infection.

207 Considering serostatus, as expected, no response was found in seronegative patients, whereas a
208 mean level of 11.2 SFU/2X10⁵ PBMCs (range, 0-211; median, 3) was found in seropositive
209 patients, with no significant difference in IgM-positive versus IgM-negative patients (mean SFU/2 x
210 10⁵ PBMCs ± SD, 3.5 ± 4.7 versus 11.6 ± 33.6; p = 0.585). In Table 2, HSV-1 responses to different
211 antigenic concentrations according to IgG serostatus are reported.

212 In order to investigate the kinetics of HSV-1-specific T-cell immunity after lung transplantation,
213 EliSPOT data were evaluated at different time points, including prior to transplantation, at 1 month
214 and at 6-month intervals posttransplantation. The pattern of HSV-1 specific cellular immune
215 response evidenced a decrease in the first months posttransplantation in comparison to
216 pretransplantation levels; this was seen with both 3.16 x 10³ and 3.16 x 10⁴ TCID₅₀/mL antigenic
217 stimuli (mean SFU/2 x 10⁵ PBMC ± SD, 2.889 ± 5.061 versus 9.5 ± 16.10, and 1.0 ± 2.0 versus 8.2
218 ±15.30, respectively), with progressive recovery of pretransplantation levels at the end of the
219 second year posttransplantation (5.444 ± 8.819 versus 4.222 ± 6.685, for 3.16 x 10³ and 3.16 x 10⁴
220 TCID₅₀/mL HSV-1 stimuli, respectively)(Figure 4, A and B). This kinetics was observed also when
221 excluding patients with HSV-1 DNA positivity on BAL specimens in concomitance with the
222 EliSPOT determinations (n=7)(Figure 4, C and D).

223 Seven lung transplant recipients (15.6%) exhibited at least one episode of HSV-1 lower respiratory
224 tract infection (as determined by molecular detection of HSV-1 DNA on BAL specimens [Costa et
225 al. 2012c]), concomitant to the available EliSPOT assays. All cases of HSV-1 infection occurred in
226 IgG-positive recipients, likely due to viral reactivation. In these patients, the level of HSV-1 cellular
227 immunity tended to be lower in comparison to patients with no HSV-1 DNA positivity, even though
228 not reaching statistical significance (mean EliSPOT values: 1.143 ± 0.5533 versus 3.967 ±0.7295;
229 p=0.1986).

230 In order to assess the impact of pulmonary events of HSV-1 replication on subsequent virus-specific
231 immunity induction, BAL determinations performed in a 6-month period prior to the available
232 EliSPOT assays were retrospectively investigated. Six patients exhibited a history of at least one
233 episode of pulmonary HSV-1 replication in this period; in these patients, HSV-1 EliSPOT response
234 tended to be higher in comparison to patients with no evidence of lower respiratory tract infection in
235 the same interval (mean SFU/2 x 10⁵ PBMC, 8.167 ± 3.229 versus 3.568 ± 0.7953, p=0.0656, using
236 3.16x10³ TCID₅₀/mL as antigenic stimulus). Moreover, no relation was found between HSV-1
237 EliSPOT responses and the occurrence of HSV-1 positivity in the subsequent 6-month period.

238 **Discussion**

239 In this study, cellular immune response to HSV-1 was evaluated by a newly developed IFN- γ
240 EliSPOT assay. Whereas Posavad and colleagues described an EliSPOT assay for HSV-2 to be used
241 in vaccine development (Posavad et al. 2011), an assay specifically designed for HSV-1 has not
242 been reported in literature and its availability could be useful for defining the role of cellular
243 immunity in the development and outcome of HSV-1 infection/reactivation, as well as in its clinical
244 and therapeutic management. Immunocompromised patients, such as transplant recipients, present
245 more frequent and severe clinical manifestations of HSV-1 infection, as well as decreased responses
246 to anti-viral treatment (Wilk et al. 2013). In most of the cases, symptomatic HSV-1 disease in adult
247 transplant recipients results from viral reactivation, particularly in the first month following
248 transplantation (Fishman 2007). Among other clinical manifestations, including disseminated
249 mucocutaneous disease, esophagitis and hepatitis, pneumonitis is described in all solid organ
250 transplant patients, but most commonly in lung and heart-lung transplant patients (Smyth et al.
251 1990). The kinetics and specificity of HSV-1 T-cell immune response during primary infection are
252 poorly known in humans. Following resolution of acute episode, specific memory T-cells are found
253 at moderate levels of 0.1-1% in immunocompetent individuals (Ouwendijk et al. 2013; Jing et al.
254 2012; Moss et al. 2012). In healthy individuals, a complex and poly-specific CD4+ and CD8+
255 response towards more than 70 different proteins has been identified, including proteins abundantly
256 present in the virion (e.g. viral envelope, tegument, capsid) and regulatory proteins (Jing et al. 2012;
257 Merindol et al. 2012; Harari et al. 2006; Jing et al. 2013). HSV-1 specific T-cells localize to sites of
258 primary, recurrent and chronic latent infections from which reactivation may occur in favoring
259 conditions, such as immunosuppression. Several studies have demonstrated that the outcome of
260 these infections depends on the efficacy of specific cellular immune response (Remakus and Sigal
261 2013; Sant and McMichael 2012; Calarota et al. 2015) and that the development of quantitative,
262 sensitive and reproducible assays for evaluation and monitoring of virus-specific T-cell response is
263 fundamental to investigate kinetics of HSV-1-specific immunity and in the clinical-therapeutic
264 management of immunocompromised patients.

265 Among methods developed for evaluating virus-specific T-cell response, the EliSPOT assay allows
266 for measurement of quantity and functionality of specific T-cells and can be used to define the
267 whole repertoire of cellular responses without MHC-restriction. EliSPOT assay detects production
268 of IFN- γ by PBMCs following stimulation with specific antigens and enumerates responsive cells
269 using anti-IFN- γ monoclonal antibodies coated onto 96-well plates and a second enzyme-

270 conjugated monoclonal antibody; spots are counted using automated EliSPOT readers with each
271 spot representing a single specific cell (Calarota et al. 2015). The most common antigenic stimuli
272 used for EliSPOT assay are pools of overlapping peptides, peptide libraries spanning entire proteins
273 or viral lysates. Given the antigenic complexity of herpesviruses which contain multiple potential
274 protein targets recognized by CD4+ cells and the dose-response curves obtained on preliminary
275 EliSPOT assays, in this study a viral lysate preparation at 3.16×10^4 and 3.16×10^3 TCID₅₀/mL
276 dilutions was used.

277 By using these two dilutions of inactivated virus, we found that 3.16×10^3 TCID₅₀/mL HSV-1
278 stimulus was significantly associated to higher level of response in comparison to 3.16×10^4
279 TCID₅₀/mL. This difference was evidenced in almost all cases with very few exceptions and
280 considering those collected in both the pre- and post-transplant settings; it could be hypothesized
281 that this is due to the degree of saturation of binding sites.

282 As regards HSV-1 specific cellular immune response in study population, there was no significant
283 difference in its level between the pre- and post- transplant period. As the higher risk (Fishman
284 2007) of HSV-1 reactivation is in the very first months (particularly up to 30 days), we evaluated
285 whether this could be attributable, at least partly, to a lower degree of virus-specific cellular
286 immune control. Although the difference was not significant, a tendency to lower levels of response
287 in the first period was found; of course, it should be taken into account the small number of
288 specimens that could have limited the statistical power of these data and the need for increasing the
289 study group. Moreover, it has to be underlined that we consider cumulative data from all the
290 specimens available for a certain period of time posttransplantation, giving the difference in number of
291 samples available at different time points.

292 Knowledge of HSV-1 serostatus, as well as of cell-mediated immune response, may be of great
293 concern to stratify patients at major risk for primary HSV-1 acquisition – either from the allograft or
294 from natural sources – after transplantation, which may be more clinically severe and prolonged due
295 to lack of immunologic memory (Wilck and Zuckerman 2013; Nichols et al. 2003). As expected, no
296 response was found in seronegative patients; on the other side, when considering seropositive
297 patients, no significant difference in the degree of virus-specific response was found between IgG-
298 and IgM-positive individuals, although values tended to be higher in patients with a serological
299 status suggesting previous infection. This observation supports the hypothesis that a higher level of
300 response is achieved following immunological boosting of memory T-cell, as already reported for

301 cytomegalovirus (Costa et al. 2014a; Rittà et al. 2015; Abate et al. 2010), Epstein-Barr virus (Rittà
302 et al. 2015) and polyomavirus BK (Costa et al. 2014b).

303 Given the occurrence of HSV-1 infection/reactivation in the lower respiratory tract and the potential
304 impact in the presence of impaired immune responses, as reported for other herpesviruses (Costa et
305 al. 2007), a study population of lung transplant recipients was chosen for clinical validation of the
306 developed HSV-1 EliSPOT assay and evaluation of kinetics of specific cellular immune response.
307 As expected, a decrease (although not significant) in the level of response in the first months
308 posttransplantation was found in comparison to pretransplantation levels, with progressive recovery
309 of these levels along a period ranging from 3 months to 2 years posttransplantation.

310 As regards HSV-1 infection in the lower respiratory tract, as evidenced by positivity to HSV-1
311 DNA on BAL specimens, all the cases occurred in IgG-positive patients, thus representing viral
312 reactivation. In terms of impact of the level of HSV-1 specific cellular immune response on viral
313 reactivation, although not statistically significant, a tendency to lower levels in the seven patients
314 with at least one episode of infection was observed, with values even lower in the presence of
315 repeated episodes. Moreover, as these data referred to concomitant evaluation of HSV-1 DNA on
316 BAL and EliSPOT assay, we also assessed the impact of pulmonary HSV-1 infection on subsequent
317 level of virus-specific cellular immune response by retrospectively investigating BAL
318 determinations in a 6-month period prior to the available EliSPOT assay. Interestingly, in patients
319 with at least one episode of pulmonary HSV-1 infection in comparison to those with no infection,
320 the degree of cellular immune response tended to be higher, thus supporting the boosting effect of
321 viral replication on the development of HSV1-specific immunity.

322 In conclusion, we have evaluated T-cell responses directed at a HSV-1 in lung transplant patients by
323 a newly developed, specific and quantitative IFN- γ EliSPOT assay and investigated the
324 immunological status and kinetics. The availability of this assay could allow for a patient's tailored
325 clinical-therapeutic management in terms of modulation of immunosuppressive therapy and use of
326 antiviral agents in the presence of HSV-1 infection/reactivation in relation to the occurrence and
327 level of virus-specific response. Further studies on larger and different populations of
328 immunocompromised and immunocompetent patients are required to define the potential of
329 quantitative evaluation of HSV-1 specific cellular immune response in different clinical settings and
330 to define cut-off levels discriminating between absence/low and strong response to be related to the
331 risk of viral infection/reactivation.

332

333

334

335 **References**

336 Abate D, Saldan A, Ficon M, Cofano S, Paciolla A, et al. (2010). Evaluation of cytomegalovirus
337 (CMV)-specific T cell immune reconstitution revealed that baseline antiviral immunity,
338 prophylaxis, or preemptive therapy but not antithymocyte globulin treatment contribute to CMV-
339 specific T cell reconstitution in kidney transplant recipients. *J Infect Dis* 202:585-594.

340 Ariotti S, Beltman JB, Chodaczek G, Hoekstra ME, van Beek AE, et al. (2012). Tissue-resident
341 memory CD8+ T cells continuously patrol skin epithelia to quickly recognize local antigen. *Proc*
342 *Natl Acad Sci U S A* 109:19739-19744.

343 Bonizzoli M, Arvia R, di Valvasone S, Liotta F, Zakrzewska K, et al. (2016). Human herpesviruses
344 respiratory infections in patients with acute respiratory distress (ARDS). *Med Microbiol Immunol*
345 205:371-379.

346 Bouza E, Giannella M, Torres MV, Catalán P, Sánchez-Carrillo C, et al. (2011). Herpes simplex
347 virus: a marker of severity in bacterial ventilator-associated pneumonia. *J Crit Care* 26:432.e1-6.

348 Bruynseels P, Jorens PG, Demey HE, Goossens H, Pattyn SR, et al. (2003). Herpes simplex virus in
349 the respiratory tract of critical care patients: a prospective study. *Lancet* 362:1536-1541.

350 Calarota SA, Aberle JH, Puchhammer-Stöckl E, Baldanti F. (2015). Approaches for monitoring of
351 non virus-specific and virus-specific T-cell response in solid organ transplantation and their clinical
352 applications. *J Clin Virol* 70:109-119.

353 Costa C, Balloco C, Sidoti F, Mantovani S, Rittà M, et al. (2014a). Evaluation of CMV-specific
354 cellular immune response by EliSPOT assay in kidney transplant patients. *J Clin Virol* 61:523-528.

355 Costa C, Curtioni A, Bergallo M, Solidoro P, Lorusso M, et al. (2011). Quantitative detection of
356 HHV-6 and HHV-7 in transbronchial biopsies from lung transplant recipients. *New Microbiol*
357 34:275-280.

358 Costa C, Delsedime L, Solidoro P, Curtoni A, Bergallo M, et al. (2012a). Herpesviruses detection
359 by quantitative real-time polymerase chain reaction in bronchoalveolar lavage and transbronchial
360 biopsy in lung transplant: viral infections and histopathological correlation. *Transplant Proc*
361 42:1270-1274.

362 Costa C, Elia M, Astegiano S, Sidoti F, Terlizzi ME, et al. (2008). Quantitative detection of
363 Epstein-Barr virus in bronchoalveolar lavage from transplant and nontransplant patients.
364 *Transplantation* 86:1389-1394.

365 Costa C, Libertucci D, Solidoro P, Sinesi F, Bergallo M, et al. (2007). Rapid shell vial culture for
366 the detection of respiratory viruses from bronchoalveolar lavage in immunocompromised patients.
367 *Panminerva Med* 49:1-6.

368 Costa C, Mantovani S, Piccighello A, Di Nauta A, Sinesi F, et al. (2014b). Evaluation of
369 polyomavirus BK cellular immune response by an EliSPOT assay and relation to viral replication in
370 kidney transplant recipients. *New Microbiol* 37:219-223.

371 Costa C, Saldan A, Sinesi F, Sidoti F, Balloco C, et al. (2012b). The lack of cytomegalovirus-
372 specific cellular immune response may contribute to the onset of organ infection and disease in lung
373 transplant recipients. *Int J Immunopathol Pharmacol* 25:1003-1009.

374 Costa C, Sidoti F, Saldan A, Sinesi F, Balloco C, et al. (2012c). Clinical impact of HSV-1 detection
375 in the lower respiratory tract from hospitalized adult patients. *Clin Microbiol Infect* 18:E305-307.

376 Daubin C, Vincent S, Vabret A, du Cheyron D, Parienti JJ, et al. (2005). Nosocomial viral
377 ventilator-associated pneumonia in the intensive care unit: a prospective cohort study. *Intensive*
378 *Care Med* 31:1116-1122.

379 De Vos N, Van Hoovels L, Vankeerberghen A, Van Vaerenbergh K, Boel A, et al. (2009).
380 Monitoring of herpes simplex virus in the lower respiratory tract of critically ill patients using real-
381 time PCR: a prospective study. *Clin Microbiol Infect* 15:358-363.

382 Engelmann I, Gottlieb J, Meier A, Sohr D, Ruhparwar A, et al. (2007). Clinical relevance of and
383 risk factors for HSV-related tracheobronchitis or pneumonia: results of an outbreak investigation.
384 *Crit Care* 11:R119.

385 Fishman JA. (2007). Infection in solid-organ transplant recipients. *N Engl J Med* 357:2601-2614.

386 Gebhardt T, Wakim LM, Eidsmo L, Reading PC, Heath WR, et al. (2009). Memory T cells in
387 nonlymphoid tissue that provide enhanced local immunity during infection with herpes simplex
388 virus. *Nat Immunol* 10:524-530.

389 Gebhardt T, Whitney PG, Zaid A, Mackay LK, Brooks AG, et al. (2011). Different patterns of
390 peripheral migration by memory CD4⁺ and CD8⁺ T cells. *Nature* 477:216-219.

391 Gooskens J, Templeton KE, Claas EC, von Bussel MJ, Smit VT, et al. (2007). Quantitative
392 detection of herpes simplex virus DNA in the lower respiratory tract. *J Med Virol* 79:597-604.

393 Harari A, Dutoit V, Cellerai C, Bart PA, Du Pasquier RA, et al. (2006). Functional signatures of
394 protective antiviral T-cell immunity in human virus infections. *Immunol Rev* 211:236-254.

395 Jing L, Haas J, Chong TM, Bruckner JJ, Dann GC, et al. (2012). Cross-presentation and genome-
396 wide screening reveal candidate T cells antigens for a herpes simplex virus type 1 vaccine. *J Clin*
397 *Invest* 122:654-673.

398 Jing L, Schiffer JT, Chong TM, Bruckner JJ, Davies DH, et al. (2013). CD4 T-cell memory to viral
399 infections of humans shows pronounced immunodominance independent of duration or viral
400 persistence. *J Virol* 87:2617-2627.

401 Khanna KM, Bonneau RH, Kinchington PR, Hendricks R. (2003). Herpes simplex virus-specific
402 memory CD8⁺ T cells are selectively activated and retained in latently infected sensory ganglia.
403 *Immunity* 18:593-603.

404 Linssen CF, Jacobs JA, Stelma FF, van Mook WN, Terporten P, et al. (2008). Herpes simplex virus
405 load in bronchoalveolar lavage fluid is related to poor outcome in critically ill patients. *Intensive*
406 *Care Med* 34:2202-2209.

407 Luyt C-E, Combes A, Deback C, Aubriot-Lorton M-H, Nieszkowska A, et al. (2007). Herpes
408 Simplex Virus lung infection in patients undergoing prolonged mechanical ventilation. *Am J Respir*
409 *Crit Care Med* 175:935-942.

410 Merindol N, Salem Fourati I, Brito RM, Grenier AJ, Charrier E, et al. (2012). Reconstitution of
411 protective immune responses against cytomegalovirus and varicella zoster virus does not require
412 disease development in pediatric recipients of umbilical cord blood transplantation. *J Immunol*
413 189:5016-5028.

414 Moss NJ, Magaret A, Laing KJ, Kask AS, Wang M, et al. (2012). Peripheral blood CD4 T-cell and
415 plasmacytoid dendritic cell (pDC) reactivity to herpes simplex virus 2 and pDC number do not
416 correlate with the clinical or virologic severity of recurrent genital herpes. *J Virol* 86:9952-9963.

417 Nichols WG, Boeckh M, Carter RA, Wald A, Corey L. (2003). Transferred herpes simplex virus
418 immunity after stem-cell transplantation: clinical implications. *J Infect Dis* 187:801-808.

419 Ong GM, Lowry K, Mahajan S, Wyatt DE, Simpson C, et al. (2004). Herpes simplex type 1
420 shedding is associated with reduced hospital survival in patients receiving assisted ventilation in a
421 tertiary referral intensive care unit. *J Med Virol* 72:121-125.

422 Ouwendijk WJD, Laing KJ, Verjans GM, Koelle DM. (2013). T-cell immunity to human
423 alphaherpesviruses. *Curr Opin Virol* 3:452-460.

424 Posavad CM, Magaret AS, Zhao L, Mueller DE, Wald A, et al. (2011). Development of an
425 interferon-gamma ELISPOT assay to detect human T cell responses to HSV-2. *Vaccine* 29:7058-
426 7066.

427 Preiser W, Doerr HW, Vogel J-U. (2003). Virology and epidemiology of oral herpesvirus
428 infections. *Med Microbiol Immunol* 192:133-136.

429 Remakus S, Sigal LJ. (2013). Memory CD8(+) T cell protection. *Adv Exp Med Biol* 785:77-86.

430 Rittà M, Costa C, Sidoti F, Balloco C, Ranghino A, et al. Pre-transplant assessment of CMV-
431 specific immune response by Elispot assay in kidney transplant recipients. (2015). *New Microbiol*
432 38:329-335.

433 Rittà M, Costa C, Sinesi F, Sidoti F, Di Nauta A, et al. (2013). Evaluation of Epstein-Barr virus-
434 specific immunologic response in solid organ transplant recipients with an enzyme-linked
435 ImmunoSpot assay. *Transplant Proc* 45:2754-2757.

436 Sant AJ, McMichael A. (2012). Revealing the role of CD4(+) T cells in viral immunity. *J Exp Med*
437 209:1391-1395.

438 Scheithauer S, Manemann AK, Krüger S, Häusler M, Krüttgen A, et al. (2010). Impact of herpes
439 simplex virus detection in the respiratory specimens of patients with suspected viral pneumonia.
440 *Infection* 38:401-405.

441 Simmoons-Smit AM, Kraan EM, Beishuizen A, Strack van Schijndel RJ, et al. (2006). Herpes
442 simplex virus type 1 and respiratory disease in critically-ill patients: Real pathogen or innocent
443 bystander? *Clin Microbiol Infect* 12:1050-1059.

444 Smyth RL, Higenbottam TW, Scott JP, Wreghitt TG, Stewart S, et al. (1990). Herpes simplex virus
445 infection in heart-lung transplant recipients. *Transplantation* 49:735-739.

446 Terlizzi ME, Bergallo M, Sidoti F, Sinesi F, Vendrame R, et al. (2009). Quantitative RT real time
447 PCR and indirect immunofluorescence for the detection of human parainfluenza virus 1, 2, 3. *J*
448 *Virology Methods* 160:172-177.

449 Tsakris A, Pitiriga VC. (2011). Herpes simplex viruses 1 and 2; in Liu D (ed): *Molecular detection*
450 *of human viral pathogens*, CRC Press, Taylor & Francis Group, Boca Raton, pp 911-922.

451 van den Brink JW, Simoons-Smit AM, Beishuizen A, Girbes AR, Strack van Schijndel RJ, et al.
452 (2004). Respiratory herpes simplex virus type 1 infection/colonization in the critically ill: marker or
453 mediator? *J Clin Virol* 30:68-72.

454 Wilck MB, Zuckerman RA. (2013). Herpes simplex virus in solid organ transplantation. *Am J*
455 *Transplant* 13(Suppl 4):121-127.

456
457
458
459
460
461
462

463 Table 1. Demographic and clinical features of study population. BAL, bronchoalveolar lavage;
 464 COPD, chronic obstructive pulmonary disease; CSA, cyclosporin A; MMF, mycophenolate mofetil;
 465 MPA, mycophenolic acid; TAC, tacrolimus; AZA, azathioprine; EVR, everolimus. Details on
 466 antiviral prophylaxis are reported in the text.

Features	
Patients, total n	95
Male/female, n	56/39
Mean age (range), years	47.2 (16-69)
Healthy seropositive individuals, n	42
Mean age (range), years	37.5 (21-49)
N. of EliSPOT determinations per patient	1
Pre-transplant patients, n	8
Mean age (range), years	47 (22-65)
N. of EliSPOT determinations per patient	1
Post-transplant patients, n	45
Mean age (range), years	47.0 (16-69)
N. of EliSPOT determinations per patient (mean, range)	3.6 (3-5)
Time of EliSPOT determinations post-transplantation (months – mean, range)	21 (1-94)
Type of lung transplant	
Monolateral	9
Bilateral	36
Underlying disease	
Cystic fibrosis	37 (50.7%)
COPD/emphysema	22 (26.0%)
Idiopathic pulmonary fibrosis	6 (8.2%)
Bronchiectasis	7 (9.6%)
Extrinsic allergic alveolitis	1 (1.4%)
Antiviral prophylaxis (in all transplant patients)	
HSV	Acyclovir
CMV	Ganciclovir or valganciclovir + CMV-Ig
Immunosuppressive regimens	
CSA + MMF	25
CSA + MPA	1
TAC + MMF	15
TAC + MPA	2
TAC + AZA	1
TAC + EVR	1
HSV-1/2 serology at baseline	
IgM+	5
IgM-	48
IgG+	48
IgG-	5

467

468

469 Table 2. HSV-1 EliSPOT responses to different antigenic concentration according to IgG serostatus
470 (mean \pm standard deviation, spot forming units [SFU]/ 2×10^5 peripheral blood mononuclear cells
471 [PBMCs]).

	HSV IgG+ (n = 48)	HSV IgG- (n = 5)	p
3.16×10^3 TCID ₅₀ /mL	5.447 \pm 1.543	0.2 \pm 0.2	0.24
3.16×10^4 TCID ₅₀ /mL	3.929 \pm 1.335	0.2 \pm 0.2	0.32

472

473

474 Figure 1. Rapid shell vial culture assay with indirect immunofluorescence using Vero cells infected
475 with (A) human Herpesvirus 1 ATCC® VR-260TM, (B) UV-inactivated HSV-1 preparation
476 (dilution 3.16×10^8 TCID₅₀/mL), and (C) UV-treated RPMI-1640 complete medium alone at 24 h
477 post-infection (Fluorescein isothiocyanate; counterstaining with Evans blue 1:10000).
478 Magnification, 25X.

479

480 Figure 2. EliSPOT assay on peripheral blood mononuclear cells from a HSV-1 IgG-positive lung
481 transplant patient stimulated with serial dilution of UV-inactivated HSV-1 preparation: (A) $3.16 \times$
482 10^4 TCID₅₀/mL, (B) 3.16×10^3 TCID₅₀/mL, (C) RPMI-1640 complete medium alone, and (D)
483 phytohemagglutinin mitogen (PHA) 1 µg/mL. Results are reported as spot forming unit
484 (SFU)/ 2×10^5 cells.

485

486 Figure 3. HSV-1 EliSPOT responses according to concentrations of HSV-1 antigenic stimulus in
487 samples from posttransplantation patients (peak value of response for each patient).

488

489 Figure 4. Kinetics of HSV-1 EliSPOT responses according to concentrations of HSV-1 antigenic
490 stimulus considering all specimens from transplant patients (A, 3.16×10^3 TCID₅₀/mL; B, $3.16 \times$
491 10^4 TCID₅₀/mL) and excluding specimens from patients with concomitant HSV-1 positivity on
492 bronchoalveolar lavage (BAL) (C, 3.16×10^3 ; D, 3.16×10^4 TCID₅₀/mL). Determinations are
493 grouped as follows: at pre-transplant (n = 42), up to 1month (n = 7), at 1-6 months (n = 13), at 6-12
494 months (n = 79), at 12-18 months (n = 53), and at 18-24 months (n = 8) post-lung transplantation
495 (LT).